


# RUTGERS

New Jersey Agricultural  
Experiment Station

## **2017 Turfgrass Proceedings**

***The New Jersey Turfgrass Association***

In Cooperation with  
Rutgers Center for Turfgrass Science  
Rutgers Cooperative Extension



# **2017 RUTGERS TURFGRASS PROCEEDINGS**

of the

## **GREEN EXPO Turf and Landscape Conference**

**December 5-7, 2017**

**Borgata Hotel**

**Atlantic City, New Jersey**

The Rutgers Turfgrass Proceedings is published yearly by the Rutgers Center for Turfgrass Science, Rutgers Cooperative Extension, and the New Jersey Agricultural Experiment Station, School of Environmental and Biological Sciences, Rutgers, The State University of New Jersey in cooperation with the New Jersey Turfgrass Association. The purpose of this document is to provide a forum for the dissemination of information and the exchange of ideas and knowledge. The proceedings provide turfgrass managers, research scientists, extension specialists, and industry personnel with opportunities to communicate with co-workers. Through this forum, these professionals also reach a more general audience, which includes the public.

This publication includes lecture notes of papers presented at the 2017 GREEN EXPO Turf and Landscape Conference. Publication of these lectures provides a readily available source of information

covering a wide range of topics and includes technical and popular presentations of importance to the turfgrass industry.

This proceedings also includes research papers that contain original research findings and reviews of selected subjects in turfgrass science. These papers are presented primarily to facilitate the timely dissemination of original turfgrass research for use by the turfgrass industry.

Special thanks are given to those who have submitted papers for this proceedings, to the New Jersey Turfgrass Association for financial assistance, and to Barbara Fitzgerald and Anne Diglio for administrative and secretarial support.

Dr. Ann Brooks Gould, Editor  
Dr. Bruce B. Clarke, Coordinator

# PRE-EMERGENCE GOOSEGRASS CONTROL WITH COMMERCIALY AVAILABLE HERBICIDES, 2017

Matthew T. Elmore and Daniel P. Tuck<sup>1</sup>

The objective of this experiment was to evaluate various commercially available herbicides for pre-emergence goosegrass (*Eleusine indica*) control.

## MATERIALS AND METHODS

This experiment was conducted at the Rutgers Horticultural Farm II in North Brunswick, NJ on a sandy loam soil with a pH of 5.5 and a history of goosegrass. A poor stand of mature perennial ryegrass (*Lolium perenne*) (turf cover ~40%) was also present. The site was mowed weekly at 1.5 inches with a reel mower and irrigated at least weekly to prevent wilt. No additional fertilizers or plant protectants were applied to the trial. Quinclorac was applied to the entire trial site in late July to control crabgrass.

Treatments (Table 1) were arranged in a randomized block design and replicated three times. The treatments were applied to 4 x 7-ft plots using a CO<sub>2</sub>-powered sprayer calibrated to apply 44 GPA through a single 9504EVS nozzle at 44 PSI. Granular treatments were applied using a shaker jar. Applications A, B, and C were made on 24 February, 13 April, and 1 June 1 2017, respectively. A 12-inch wide, non-treated buffer strip was maintained between each plot providing a 3 x 7-ft treated area.

Goosegrass control and perennial ryegrass injury was evaluated visually on a 0 (no injury or control) to 100% (complete control) scale relative to the non-treated control. Goosegrass cover was moderate (~30%) in the non-treated control plots at the final evaluation in September. Data were analyzed subjected to ANOVA in ARM (v2017) and Fisher's Protected LSD ( $p \leq 0.05$ ) was used to separate means.

## RESULTS

No perennial ryegrass injury was observed at any time during the experiment (data not presented).

Treatments (Table 2) that provided similar goosegrass control at 15 weeks after the C application included: 1) sequential applications of prodiamine; 2) sequential applications of oxadiazon; 3) a single oxadiazon application in April; 4) prodiamine applied singly in February; and 5) sequential applications of dithiopyr in April/June. No treatment provided >90% goosegrass control. Single applications of dithiopyr provided poor (<50%) goosegrass control.

---

<sup>1</sup>Assistant Extension Specialist in Weed Science and Field Researcher IV, respectively, New Jersey Agricultural Experiment Station, School of Environmental and Biological Sciences, Rutgers, The State University of New Jersey, New Brunswick, NJ 08901-8520.

Table 1. Herbicide treatments applied for pre-emergence goosegrass (*Eleusine indica*) control at the Rutgers Horticultural Research Farm II, North Brunswick, NJ. Applications A, B, and C were made on 24 February, 13 April, and 1 June, 2017.

Treatment	Product	Active Ingredient	Product Rate (per acre)	Active Ingredient Rate (lb per A)	Application Code
1	Non-treated	–	–	–	–
2	Dimension 2EW	dithiopyr	32 fl oz	0.5	A
3	Dimension 2EW	dithiopyr	32 fl oz	0.5	B
4	Ronstar 2G	oxadiazon	87 lb	1.7	B
5	Echelon	prodiamine + sulfentrazone	12 fl oz	0.22 + 0.11	B <i>fb</i> <sup>1</sup> C
6	Barricade 65WG	prodiamine	1.54 lb	1.0	A
7	Barricade 65WG	prodiamine	1.54 lb	1.0	B
8	Dimension 2EW	dithiopyr	16 fl oz	0.25	A <i>fb</i> C
9	Barricade 65WG	prodiamine	0.77 lb	0.5	A <i>fb</i> C
10	Ronstar 2G	oxadiazon	75 lb	1.5	B <i>fb</i> C
11	Dimension 2EW	dithiopyr	16 fl oz	0.25	B <i>fb</i> C
12	Barricade 65WG	prodiamine	0.77 lb	0.5	B <i>fb</i> C
13	Ronstar 2G	oxadiazon	44 lb	0.65	B <i>fb</i> C

<sup>1</sup> *fb* = followed by

Table 2. Goosegrass control from pre-emergence herbicide applications in North Brunswick, NJ. Applications A, B, and C were made on 24 February, 13 April, and 1 June, 2017.

Treatment	Product	Application Code	Goosegrass Control (%) <sup>1</sup>	
			17 Aug. 25 WA-A <sup>2</sup> 11 WA-C <sup>3</sup>	18 Sept. 29 WA-A 15 WA-C
1	Non-treated	–	0 cd	0 e
2	dithiopyr (0.5 lb)	A	2 d	3 e
3	dithiopyr (0.5 lb)	B	14 bcd	48 cd
4	oxadiazon (1.7 lb)	B	38 abc	63 a-d
5	prodiamine + sulfentrazone	B <i>fb</i> <sup>4</sup> C	35 abc	55 cd
6	prodiamine (1.0 lb)	A	69 a	85 a
7	prodiamine (1.0 lb)	B	37 abc	60 bcd
8	dithiopyr (0.5 lb)	A <i>fb</i> C	28 a-d	45 d
9	prodiamine (0.5 lb)	A <i>fb</i> C	64 a	87 a
10	oxadiazon (1.5 lb)	B <i>fb</i> C	52 ab	83 ab
11	dithiopyr (0.25 lb)	B <i>fb</i> C	55 ab	70 abc
12	prodiamine (0.5 lb)	B <i>fb</i> C	35 abc	72 abc
13	oxadiazon (0.65 lb)	B <i>fb</i> C	59 a	80 ab
LSD at 5% =			29-40	25

<sup>1</sup> Goosegrass control evaluated on a 0 to 100% scale, where 0 = no control and 100 = complete control relative to the non-treated control. Means followed by the same letter are not significantly different according to Fisher's Protected LSD test ( $p \leq 0.05$ )

<sup>2</sup> WA-A = weeks after application A

<sup>3</sup> WA-C = weeks after application C

<sup>4</sup> *fb* = followed by